COVID-19 Patient PPE



Affirming life at every step of your journey with illness and loss.

For any Tier category: If clinician determines use of N95 is appropriate for any visit, clinician should wear N95.

***If unable to fit with N95, a KN95 or procedure mask are acceptable according to CDC emergency recommendations. PAPR units are also available at the Telecare Center and the Care Center. ***

Tier	Tier 1	Tier 2	Tier 3	Tier 4	Tier 5
Definition	COVID+ Patient	Close Contact with COVID+ Case	Positive Screened Patient	Positive Screened Family/Caregiver in Home	Patients Screen No for COVID Exposure
Home Emergency Plan	Whenever possible, patient to:	Patient/family should quarantine at home with	Patient should quarantine at home with minimal to no	Family or CG should quarantine from patient to	Routine infection control precautions
	1) Wear N95 or K95 mask at all times	minimal to no visitors	visitors	the best of their ability	
	Quarantine to individual room with its own bathroom	Sick individuals should wear mask at all times			Refer to Relias course Infection Control: The Basics
	3) Use disposable utensils 4) Use independent trash can				REL-ALL-0-ICBASIC
Staffing	1) Visits by all members of IDT as appropriate and with appropriate PPE 2) Telehealth visits when	1) Visits by all members of IDT as appropriate and with appropriate PPE	1)Visits by all members of IDT as appropriate and with appropriate PPE	Visits by all members of IDT as appropriate and with appropriate PPE	Routine staffing- refer to visit protocols
	possible	2) Telehealth visits when possible			

	All PPE except shield or goggles: one-time use *N95 with or without valve OR K95 or procedure mask Gown Shoe covers Gloves	All PPE except shield or goggles: one-time use *N95 with or without valve OR K95 or procedure mask Gown Shoe covers Gloves	All PPE except shield or goggles: one-time use *N95 with or without valve OR K95 or procedure mask Gown Shoe covers Gloves	All PPE except shield or goggles: one-time use *N95 with or without valve OR K95 or procedure mask Gown Shoe covers Gloves	All PPE except shield or goggles: one-time use *N95 with or without valve OR K95 or procedure mask Gown Shoe covers Gloves
PPE	Request patient and caregiver wear a mask during visit. Provide cloth mask as needed *N95 with valve worn with a procedure mask over may be required in facilities. Please refer to individual facility policies for PPE requirements if different from TRU policy.	*N95 with valve worn with a procedure mask over may be required in facilities. Please refer to individual facility policies for PPE requirements if different from TRU policy.	*N95 with valve worn with a procedure mask over may be required in facilities. Please refer to individual facility policies for PPE requirements if different from TRU policy.	*N95 with valve worn with a procedure mask over may be required in facilities. Please refer to individual facility policies for PPE requirements if different from TRU policy.	**For face shields and goggles all tiers: Wiped down and reused: use disinfecting wipes such as sanicloths to clean shield or goggles after each visit. Discard when visible signs of wear present or damaged. *N95 with valve worn with a procedure mask over may be required in facilities. Please refer to individual facility policies for PPE requirements if different from TRU policy.

^{*}We continue to check for N95 availability in multiple sizes through our trusted vendors, but they are not available at this time. Clinical Managers will work with individuals on the appropriate PPE given the current conditions. Availability of PPE is subject to change as supply chains change.

N95 for nebulizing procedures or within 2 hours of procedure, CPAP, BiPap or trilogy ventilator

All staff to wear a mask and protective eyewear at all visits.